

Molecular Characterization of a Rigid Rod-Shaped Virus Isolated from Frangipani (*Plumeria* sp.) Showing Mosaic Symptom in Taiwan

Fery Abdul Choliq^{1*}, Tsang-Hai Chen², Liliek Sulistyowati³

^{1,3}Department of Plant Protection, Faculty of Agriculture, University of Brawijaya, Malang, Indonesia

²Department of Plant Medicine, National Pingtung University of Science and Technology, Pingtung, Taiwan

Abstract

Frangipani is an important succulent plant around the worlds and also in Taiwan, for example, *Plumeria rubra* is widely grown as a popular ornamental tree in parks and landscaped establishments in Taiwan. Recently, a new disease in frangipani with mosaic and distortion symptoms was found in Taiwan. No viruses caused frangipani disease has been reported in Taiwan and the references about frangipani disease are still limited and only *Frangipani mosaic virus* (FrMV) was found. In this study, the molecular properties of a virus isolated from symptomatic frangipani in south Taiwan, such as Pingtung, Kauhsiung and Tainan were investigated. The virus with rod-shaped particles of 300 nm long and 18 nm in diameter was examined inside diseased leaves by electron microscopy. The purified virus particles showed the typical UV spectrum of tobamoviruses with A_{260}/A_{280} value of 1.29 and maximum and minimum absorption at 260 nm and 249 nm, respectively. The molecular weight of 19.5 kDa as the size of coat protein of tobamoviruses was estimated by sodium dodecyl sulfate-polyacrylamide gel (SDS-PAGE). Furthermore, the degenerate primers for tobamoviruses were used to amplify 568 bp and 400 bp of the DNA fragments in RT-PCR and nested PCR, respectively. Based on these results, it was confirmed that the rigid rod-shaped virus isolated from mosaic symptom of frangipani leaves is an isolate of FrMV, belonging to the genus *Tobamovirus*. This is the first report that FrMV infecting *Plumeria* sp. in Taiwan.

Keywords: Frangipani plant, FrMV, mosaic disease, *Tobamovirus*.

INTRODUCTION

Frangipani (*Plumeria* sp.) is a small group of plant species native to tropical countries [1]. Frangipani is an important succulent plant around the worlds and some species are valuable sources as medicines, insecticides, fibers, and rubber [2]. Frangipani plants are also important in Taiwan.

For the example is *P. rubra* which widely grown as a popular ornamental tree in parks and landscaped establishments in Taiwan. It bears beautiful, big flowers of various colours and sizes that predominate especially during the summer [3].

Unfortunately, there are factors which inhibit frangipani growth, i.e. pest and pathogen attacked. One of pathogens which could attack frangipani plant is virus. Preliminary surveys in the fields of Pingtung, Kauhsiung and Tainan County showed that some frangipani plants are indicating attacked by viruses based on foliar symptoms about 30% severity (unpublished data, 2012). This condition encourages the research about molecular characterization

(electron microscopy, virus purification, and RT-PCR,) of unknown reported mosaic disease of *Plumeria* sp. in Taiwan. The objective of this study was to identify the virus which attacked *Plumeria* sp.

MATERIALS AND METHODS

Virus Source and Isolation

Frangipani plants showing symptom caused virus including mosaic and leaf distortion were collected in Pingtung County, Taiwan by random sampling method from 100 plants as sampel. Symptoms were recorded (photographed) at Plant Protection Laboratory, Department of Plant Medicine, National Pingtung University of Science and Technology (NPUST) Taiwan. Then, virus was isolated by mechanical inoculation from the infected frangipani leaf to indicator plant, *Chenopodium quinoa* Willd. and then local lesions of *C. quinoa* were collected and maintained on *C. quinoa* leaves by three passages [4]. Plants' leaves were dusted with carborundum (400 mesh) and 0.1 M phosphate buffer (PB) containing virus-stabilizing additives of 0.1 % 2-mercaptoethanol, pH 7.0 was used [5]. The virus was subsequently multiplied on *C. quinoa* Plants because it's highly susceptible and easy to growth and maintained in the laboratory at room temperature with supplemented fluorescence light providing a photoperiod at

* Correspondence author:

Fery Abdul Choliq

Email : feryac@ub.ac.id

Address : Department of Plant Protection, Faculty of Agriculture, University of Brawijaya, Jl. Veteran Malang, 65145

12L:12D. The virus isolate was temporary designated as Frangipani-Taiwan 1 (Fr-T1).

Electron Microscopy (Negative Staining)

Virus particles from *C. quinoa* and frangipani leaf extracts were one drop floated onto electron microscopy Formvar-fronted, carbon coated, 200 mesh copper grids and incubated for five minutes. Then, the grids were washed with 10 drops of distilled water, negatively stained with 6 drops of 2% aqueous uranyl acetate (pH 5.0) and incubated for five minutes. Then, all the fluid was removed by touching the edge of the grid carefully with a filter paper [6]. Then air dried before positioning the grid in a suitable container (grid storage box) and examination with Hitachi 7500 electron microscope. Particle size was determined by measuring the average of length and diameter of 100 virions.

The virus isolate Fr-T1 was partially purified according to the method by Dijkstra and de Jager [7] with slight modification as follows: infected *C. quinoa* leaves with local lesion symptoms were kept at deep freezer (-80°C) homogenized with one volume (v/w) 0.5 M phosphate buffer, pH 8.5, containing 0.01 M Na-EDTA. Then strained through 2 layer cheesecloth. After filtering through cheesecloth, the extract was clarified by adding 12% chloroform (4°C) and stirred for 2 minutes. Then continued by centrifuged at 8000 rpm for 20 minutes (R14A rotor, Hitachi CR21G, Japan). Collect the supernatant and layered on the top with 5 ml 20% sucrose in the bottom. Then centrifuged at 28,000 rpm for 2.5 hour (P70T rotor, Hitachi CP90WX, Japan).

The pellet was resuspended in borate buffer pH 8.2 containing 0.01 M Na-EDTA. Then centrifuge at 3,000 rpm for 10 minutes (R20A2, Hitachi CR21G, Japan). Collect the supernatant and centrifuged in swing-bucket rotor at 38,000 rpm for 23 hours (RPS40T-1180 rotor, Hitachi CP90WX) in 30% Cs₂SO₄ by density gradient centrifugation (Swing-bucket rotor). Dialysed the virus band with phosphate buffer 0.01 M, pH 7.0 and stirred for 16 hours at 4°C. Then, the purified virus was test by measuring wavelength absorbance at range 220-320 nm (Hitachi U-2001 spectrophotometer). Virus concentration (c in mg.mL⁻¹) were calculated by formula [7]:

$$c = A_{260}/E \times \text{dilution factor}$$

Determination of the Molecular Weight of Viral Coat Protein by SDS-Polyacrylamide Gel Electrophoresis (SDS-PAGE)

Healthy *C. quinoa* leaf, infected *C. quinoa* leaf, healthy frangipani leaf, field mosaic frangipani leaf, and purified virus isolate Fr-T1 were homogenized with 1 : 4 (w/v) 0.5 M phosphate buffer, pH 7.0. Then centrifuged the samples on centrifuge 10,000 rpm for 10 min and collected the supernatant. The extracts were analyzed by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) with 12% running gel and 5% stacking gel. Mix the samples with dye on comparison 4:1 in microtube. Then heat the microtube on dry bath incubator (Violet Bioscience, Inc.) at 100°C for 3 min. Load Protech (Prestained Protein Marker) 5µl as marker and 10 µl of each samples into the wells in the stacking gel. Start the electrophoresis with 70 voltage (V) during 30 minutes and continue with 120 V during 70 min. After finish running, the gel were stained with Coomassie Blue and shake gently on Orbital Shaker PSU-10i (Grant-Bio) with 70 rpm for 1 h. Replace the Coomassie Blue with destaining buffer (100 ml methanol, glacial acetic 70 ml, add ddH₂O to 1,000 ml) and shake gently on Orbital Shaker PSU-10i (Grant-Bio) with 70 rpm for 24 h. Protein bands are stained and the molecular weight of the viral coat protein was determined by comparing migration of the viral protein in the gel with that of marker protein and run in parallel lanes [7].

Amplification and Analysis of Nucleic Acid Sequences

Extraction of Total RNA

Purified virus of virus isolate Fr-T1, healthy *Nicotina benthamiana* and *C. quinoa* leaf, *Tobacco mosaic virus* (TMV) infected tomato and *N. tabacum* samples and *Odontoglossum ringspot virus* (ORSV) were prepared. Each sample take 5 µl and use RNA extraction kit (Direct-zolTM RNA miniprep) to extract total RNA from the each sample according to the manufacturer's instructions (The Epigenetics Company).

Reverse Transcription-Polymerase Chain Reaction (RT-PCR)

One-tube of RT-PCR reactions (25 µl) were performed. A final concentration of 1 µM for each tobamoviruses degenerate primers TobRT up1 (5'-GARTAYSCIGCIYTICARAC-3') and TobRT do2 (5' BGCTCRAARTTCCA-3'), was used. The cycling profile was as follows : first step at 43°C for 60 min, second step at 50°C for 2 min, third

step at 94°C for 4 min; five cycles segmented into step (a) 30 s at 95°C, step (b) 30 s at 43°C, step (c) 15 s at 72°C; 35 cycles segmented into step (a) 30 s at 95°C, step (b) 30 s at 46°C, step (c) 15 s at 72°C, followed by a final extension step at 72°C for 2 min [8]. Amplification was carried out in Px2 Thermal cycler (Thermo Electron Corporation).

Nested Polymerase Chain Reaction (PCR)

Nested PCR reactions (20 µl) were performed using 1 µl of the first RT-PCR product mixed with PCR master kit (GeneMark, Taiwan) and 1 µM of each degenerate primer TobN up3 (5'-GGCGYTGCARACIATHGTITAYCA-3'), TobN do4 (5' GTRTTICCIATRAAIGTIGTIACRTC-3') and TobN do4G (5' GCCGATRAAGGTGGTGACRTC-3'). The cycling profile consisted of a denaturing step at 95°C for 3 min, two cycles segmented into step (a) 20 s at 95°C, step (b) 15 s at 51°C, step (c) 5 s at 72°C; 26 cycles segmented into step (a) 20 s at 95°C, step (b) 15 s at 61°C, step (c) 5 s at 72°C, followed by a final extension step at 72°C for 2 min [8]. Amplification were carried out in Px2 Thermal cycler (Thermo Electron Corporation).

Electrophoresis Analysis

Electrophoresis analysis was used to ascertaining DNA product was amplified by RT-PCR and nested PCR. Get 1 µl loading dye with 5 µl PCR product uniformly mixing. Load 2 µl marker and each sample inside the horizontal electrophoresis analysis (1.5% agarose + 0.5 µl Save view DNA stain). Running on electrophoresis machine (Major Science®) with 120 V electricity for 20 min. Then, visualized under UV light and estimate the amplified product.

RESULT AND DISCUSSION

Virus Isolation

Frangipani plants exhibiting virus-like disease collected in Pingtung County (South Taiwan) which showed mosaic and leaf distortion on young leaf of frangipani plants (Fig. 1). Frangipani plants exhibiting virus-like disease was indicated on several city (Pingtung, Kauhsiung and Tainan). The symptom is similar according to previous study [9] noted that the leaves of frangipani infected by virus was showed chlorotic ringspots or mosaic and were often distorted. In addition, the virus induced chlorotic ringspot and mosaic, and often distortion on leaves of frangipani [10]. Mechanical inoculation to indicator plants

(*Chenopodium quinoa*) and isolation by three time single lesion showing local lesion on leaves (Fig. 2).



Figure 1. Mosaic symptom of frangipani leaves on the field.

Mechanical inoculation to indicator plant (*C. quinoa*) and maintenance by three time single lesion showing local lesion (Fig. 2). The local lesion were produced on the inoculated leaves of *C. Quinoa* [9]. The virus induces chlorotic ringspots or mosaic, and oftendistortion on leaves of frangipani, and severe mosaic symptoms occur on leaves of virus-infected *Nicotiana benthamiana* [10].



Figure 2. The local lesions caused by viru isolate Fr-T1 on *Chenopodium quinoa* leaf

Electron Microscopy

Rigid, rod-shaped particles with average length of 250-300 nm x 18 nm were seen in negatively stained preparations from infected frangipani leaves, infected *C. quinoa* leaves after three times single lesion and purified virus isolate Fr-T1 (Fig. 3). Electron microscopy test were continued for counting the average size (length and diameter) of 100 virions.

Rigid, rod-shaped particles with average length of 300 nm x 18 nm were seen in negatively stained preparations has constant results from infected frangipani leaves and purified virus isolate (Fig. 3). Particle morphology which has rod-shaped, usually straight with the size about 300 nm long and 18 nm in diameter are characteristic similar to FrMV [7,11,12].




Figure 3. Electron micrograph of negatively stained (2% uranyl acetate) virus particles from crude sap of frangipani leaves showing mosaic symptom (Bar=500nm).

The virus were checked the absorbance value by spectrophotometer and showed the curve spectrum rises rather slowly as the wavelength decreases from 300 to 250 nm, and then rapidly as the wavelength decreases below 250 nm. The purified virus showed the typical UV spectrum of nucleoprotein with $A_{260/280}$ value is 1.29 and maximum and minimum

absorption at 260 nm and 249 nm, respectively (Fig. 4). There is also evidence that at least at wavelength longer than 250 nm, the virus protein and RNA is partially protected from damage by UV [13].

The molecular weight of coat protein of the virus was estimated 19.5 kDa by electrophoresis in sodium dodecyl sulfate polyacrylamide gel (SDS-PAGE) (Fig. 5). Lane 1 and 3 are healthy leaf samples (*C. quinoa* and field mosaic frangipani leaf) that did not show band on 19.5 kDa, but infected leaf samples (lane 2 and 4) were showing band with identical molecular weight compare to purified virus isolate Fr-T1 (19.5 kDa). This data is slightly different from known molecular weight of tobamoviruses genus, such as the coat protein of TMV has molecular weight of 17–18 kDa [14,15]. The slight difference of coat protein molecular weight between virus isolate Fr-T1 and other tobamoviruses may be as a result when draw horizontal line for comparing marker and samples bands on gel.




Figure 4. The ultraviolet absorption spectrum of purified preparation of virus isolate Fr-T1.




Figure 5. The determination of viral coat proteins molecular weight of virus isolate Fr-T1 by SDS-polyacrylamide gel electrophoresis (SDS-PAGE). (M : Marker, 1 : Healthy *Chenopodium quinoa* leaf, 2 : Infected *Chenopodium quinoa* leaf, 3 : Healthy frangipani leaf, 4 : Field mosaic frangipani leaf, 5 : purified virus isolate Fr-T1).




Figure 6. Agarose gel electrophoresis analysis of RT-PCR and Nested PCR products obtained from different tobamovirus isolates. (M : Marker, Fr-T1 : *Frangipani virus* isolate Taiwan 1, TMV-to : *Tobacco mosaic virus* infected tomato, TMV-nb : *Tobacco mosaic virus* infected *Nicotiana tabacum*, YCI : Yocai virus isolate from company, H nb : Healthy *Nicotiana tabacum*, H cq : Healthy *Chenopodium quinoa*, NC : Negative control, ORSV : *Odontoglossum ringspot virus*).

Agarose gel electrophoresis analysis of RT-PCR (Fig. 6 left) and Nested PCR (Fig. 6 right) products obtained from different tobamovirus isolates were conducted for detection and partial characterization of unknown species. The virus isolate Fr-T1 could be amplified by RT-PCR using degenerate primers TobRT up1 and TobRT do2 and resulted 568 bp. This length are similar comparing to other tobamoviruses isolates (TMV-to, TMV-nb and ORSV) [8].

Nested PCR assays using Fr-TI isolate from RT-PCR product yielded about 400 bp and this amplification products is similar as expected for all tobamovirus isolates tested (TMV-to, TMV-nb and ORSV), but not shown on control (Healthy *Nicotina benthamiana* and *C. quinoa*) (Fig. 7 right). Degenerate primers for detection of tobamoviruses by RT-PCR can be more efficient for the amplification of most members of a gene family, also allowing for the detection of new and unidentified virus species, followed by a nested PCR amplification that increased specificity and sensitivity of detection [8].

CONCLUSION

Based on these results, it was confirmed that the rigid rod-shaped virus isolated from mosaic symptom of frangipani leaves has length of 300 nm x 18 nm and could be identified as FrMV, belongs to *Tobamovirus* upon from its molecular (SDS-PAGE, RT-PCR, Nested PCR and nucleotide

sequence comparison) characterizations. This is the first report that FrMV infecting frangipani (*Plumeria* sp.) in Taiwan.

REFERENCES

- [1] Eggli, U. 2002. Illustrated handbook of succulent plants (Dicotyledons). Springer. Switzerland.
- [2] Tao, L.P., A.J.M. Leeuwenberg, D.J. Middleton. 1995. Apocynaceae. *Flora of China* 16. 143-188.
- [3] Chung, W.H., C.P. Abe, Y. Yamaoka, T.W. Haung, M. Kakishima. 2005. The first report of *Plumeria* (Frangipani) rust disease caused by *Coleosporium plumeriae* in Taiwan. BSPP. New Disease Reports 11: 16.
- [4] Chen, T.H., Y.T. Lu. 1995. Partial characterization and ecology of bamboo mosaic Potexvirus from bamboos in Taiwan. *Plant Pathol. Bull.* 4. 83-90.
- [5] Valiunas, D., M. Samuitiene, M. Navalinskiene, R.E. Davis. 2008. Identification of viral and phytoplasmal agents causing disease in *Gaillardia* Foug. *Plants in Lithuania. Agro. Res.* 6. 109-118.
- [6] Hajibagheri, M.A. 1999. Electron microscopy methods and protocols. Humana Press. Totowa. New Jersey.
- [7] Dijkstra, J., C.P. de Jager. 1998. Practical plant virology: Protocols and exercises. Springer-Verlag. Berlin.

- [8] Doivas, C.I., K. Efthimiou, N.I. Katis. 2004. Generic detection and differentiation of tobamoviruses by spot nested RT-PCR-RFLP using dl-containing primers along with homologous dG-containing primers. *J. Virol. Methods*. 117. 137-144.
- [9] Francki, R.I.B., M. Zaitlin, C.J. Grivel. 1971. An unusual strain of tobacco mosaic virus from *Plumeria acutifolia*. *Aust. J. Biol. Sci.* 24. 815-818.
- [10] Lim, M.A., J.S. Hong, Y.S. Song, K.H. Ryu. 2010. The complete genome sequence and genome structure of *Frangipani mosaic virus*. *Arch. Virol.* 155. 1543–1546.
- [11] VIDE Database. 2011. Plant Viruses Online: *Frangipani mosaic virus*. Available at: <http://sdb.im.ac.cn/vide/descr345.htm>.
- [12] Varma, A., A.J. Gibbs. 1970. Frangipani mosaic virus. Available at: <http://www.dpvweb.net/dpv/showdpv.php?dpvno=196>.
- [13] Kleczkowski, A., A.D. McLaren. 1967. Inactivation of Infectivity of RNA of Tobacco Mosaic Virus during ultraviolet irradiation of the whole virus at two wavelengths. *J. Gen. Virol.* 1. 441-448.
- [14] Astier, S., J. Albouy, Y. Maury, C. Robaglia, H. Lecoq. 2007. Principles of plant virology: genome, pathogenicity, virus ecology. Science Publisher. USA.
- [15] Hull, R. 2009. Comparative plant virology, 2nd Ed. Academic Press. UK.