Microbial Community Composition of Two Environmentally Conserved Estuaries in the Midorikawa River and Shirakawa River

Tran Thanh Liem^{1*}, Mitsuaki Nakano¹, Hiroto Ohta¹, Takuro Niidome¹, Tatsuya Masuda²,
Kiyoshi Takikawa³, Shigeru Morimura¹

¹Graduate School of Science and Technology, Kumamoto University, Kumamoto City, Japan

²Priority Organization for Innovation and Excellence, Kumamoto University, Kumamoto City, Japan

³Center for Marine Environment Studies, Kumamoto University, Kumamoto City, Japan

Abstract

To provide a general overview of the microbial communities in environmentally conserved estuaries, the top 5 cm of sediment was sampled from the sandy estuary of the Shirakawa River and from the muddy estuary of the Midorikawa River. Higher amounts of organic matter were detected in the Midorikawa estuary sample than in the Shirakawa estuary sample. Measurement of redox potential revealed that the Shirakawa estuary was aerobic and the Midorikawa estuary was much less aerobic. Clone analysis was performed by targeting partial 16S rRNA gene sequences and using extracted DNA from the samples as a template. Various bacteria were detected, among which *Gammaproteobacteria* was dominant at both estuaries. Unclassified clones were detected in the *Gammaproteobacteria* group, mainly among samples from the Midorikawa estuary. Other detected bacterial groups were *Alphaproteobacteria*, *Deltaproteobacteria*, *Chloroflexi*, *Actinobacteria*, and *Bacteroidetes*. All the *Deltaproteobacteria* clones were anaerobic sulfate-reducing bacteria. Those aerobic and anaerobic bacteria coexisted in the top 5 cm of the estuary sediments indicating the surface layer have active sulfur and carbon cycle. Abundance of aerobic *Gammaproteobacteria* may be an indicator for conserved estuaries.

Keywords: conserved environment, clone analysis, estuary, microbial community, 16S rRNA gene.

INTRODUCTION

In recent years, environmental conditions in the Ariake Sea, which contains roughly 40% of Japan's tidal flats, have worsened, leading to problems such as red tides and reduced fishery output. Tidal flats are extremely valuable ecosystems, serving as a habitat and playing diverse roles in processes such as water purification and biological production. Microorganisms are a key part of the tidal flat system. In sandy tidal flats aerobic bacteria utilize and degrade organic compounds; however, in muddy tidal flats only the surface, not lower layers of sediment, is exposed to oxygen. Under these anaerobic conditions, hydrogen sulfide is easily generated by sulfate-reducing bacteria (SRB), producing a dark sediment color [1]. Although muddy tidal flats that have been environmentally conserved have nested holes populated by aerobic macrobenthic organisms, demonstrating that a muddy ecosystem itself is

not problematic, an imbalance in the metabolisms of anaerobic microorganisms will lead to harmful accumulation of hydrogen sulfide.

Changes have been reported in the microbial community of marine sediment following environmental contamination with pollutants such as oil [2-4] or heavy metals [5,6], but only a few studies have observed microbial communities under environmentally conserved conditions [7,8]. In the present study, two estuaries connecting to the Ariake Sea were sampled: the Shirakawa River estuary with sandy sediment and the Midorikawa River estuary with muddy sediment. Sites were chosen due to their high numbers of nested holes, the presence of salt-tolerant plants growing at higher water marks, and their conserved environmental conditions. Microbial communities were analyzed by targeting partial 16S rRNA gene sequences to provide a general overview of the microbial communities in environmentally conserved tidal flat sediment.

* Correspondence author:

Tran Thanh Liem

Email : liemmt2003@yahoo.com

Address : Graduate School of Science and Technology,
Kumamoto University, 2-39-1 Kurokami Chuo-ku,
Kumamoto City, Kumamoto 860-8555, Japan

MATERIALS AND METHODS

Sampling

Estuary sediment were sampled using sterilized spatula from the left bank of the Shirakawa River (N32°46'38.9", E130°36'33.6") and the right bank of the Midorikawa River (N32°42'48.3", E130°37'21.2") on May 22, 2008 and January 27, 2009. Both rivers are located in coastal areas bordering the Ariake Sea in Kumamoto Prefecture, Japan (Fig. 1). All sediment samples were collected to a depth of 5 cm at low tide. At each sampling site, samples were collected at both the high tide position and the low tide position, as shown in Fig. 2. All samples were kept at -20°C until used for analysis.

Figure 1. Sampling site Location of sampling sites in the Shirakawa and Midorikawa River estuaries connecting to the Ariake Sea

DNA Extraction

Frozen samples were thawed on ice and centrifuged at $10,000 \times g$ at 4°C for 15 min. The precipitate was washed in sterilized phosphate

buffered saline solution and was centrifuged again at the same settings. The supernatant was removed. The precipitate was mixed well and 0.5 g of the sample was used to extract DNA with the Fast DNA SPIN Kit for Soil (MP Biomedicals, Solon, OH) according to the manufacturer's protocol. DNA concentrations were measured at wavelength of 260nm (Beckman DU 530 UV-VIS Spectrophotometer).

Determination of Microbial Community

The extracted DNA was used as a template for polymerase chain reaction (PCR) targeting partial 16S rRNA gene sequences. The primer set 27F (5'-AGA GTT TGA TCC TGG CTC AG-3') and 518R (5'-GTA TTA CCG CGG CTG CTG G-3') and AmpliTaq Gold (Applied Biosystems, Carlsbad, CA) were used. After preheating at 95°C for 5 min, 25 cycles were performed of heating at 95°C for 1 min, annealing at 50°C for 1 min, and extension at 72°C for 2 min. PCR products were purified by an Ultra Clean PCR Clean-up Kit (MO-BIO, Carlsbad, CA) according to the manufacturer's protocol. Purified PCR products were ligated with pT7Blue vector (Novagen, Darmstadt, Germany), using a Ligation Mix Kit (Takara, Kyoto, Japan) according to the manufacturer's protocol. Then, 100 μ L of competent *Escherichia coli* DH5 α cells (Takara, Kyoto, Japan) was transformed with 10 μ L of ligation mixture as mentioned in manufacturer's protocol. After colonies formed on an LB-ampicillin plate containing X-gal, white colonies were selected and plasmids were extracted using the Wizard SV Minipreps DNA Purification System (Promega, Madison, WI). Extracted plasmids were digested by *Eco*RI and *Pst*I (Takara, Kyoto, Japan) to determine the size of the inserted DNA. Sequence analysis of the inserted DNA was completed by Takara Co., Ltd. by premixed sequencing with M13 (-47) primer (5'-CGC CAG GGT TTT CCC AGT CAC GAC-3').

Figure 2. Location of sampling points at each estuary

Data from the analysis were compared to the NCBI database by a BLAST search to identify obtained clones [9]. 16S rRNA nucleic acid sequences were aligned using ClustalX 1.8 software [10]. Phylogenetic trees were constructed using MEGA 4 software [11]. In which, nucleotide sequences were used to build phylogenetic trees using Distance-based method that measures the pairwise distance between genes. More detail, we chose Neighbor-Joining method in Bootstrap test of Phylogeny option from Phylogeny menu, selected Pairwise Deletion for Gaps/Missing Data to retain all sites containing alignment gaps and missing information before start computing.

Environmental Parameters Analysis

Hydrogen ion and NaCl concentration were measured using a handheld pH meter and surface salinity meter (SSM14-P and WM-22EP; TOADKK, Tokyo, Japan). Redox potential (ORP) was measured using a portable ORP meter (RM-20P; TOADKK) and distribution of particle size was analyzed using a laser diffraction particle size analyzer (LA-920; HORIBA, Kyoto, Japan). Organic matter was determined by measurement of residual weight after heating according to the Japanese Standard Method, JIS-K0102 [12]. The

partial 16S rRNA gene sequences obtained in this work have been deposited in the DNA Data Bank of Japan (DDBJ) under accession numbers AB853487–AB853725.

RESULTS AND DISCUSSION

Environmental Conditions at Sampling Sites

Environmental conditions at the Midorikawa and Shirakawa estuaries are summarized in Table 1. Generally, pH was in the range 7.5 and 8.0. Because river water and seawater were mixed at the sites, NaCl concentration was lower than 3.5%, the average concentration of seawater, due to dilution by fresh water from the rivers. ORP values were positive for the Shirakawa estuary and negative for the Midorikawa estuary, suggesting that their conditions were aerobic and slightly anaerobic, respectively. Organic content in the sediment at the Midorikawa estuary was higher than that at the Shirakawa estuary. Median particle size data and mud content analysis suggested that the Shirakawa estuary was sandy and the Midorikawa estuary was muddy. Taken together, these results confirmed that ORP was low and organic content was high under conditions of low median particle size and high mud content.

Table 1. Environmental Conditions at Shirakawa and Midorikawa River Estuary

Season	Location of estuaries	Tide level	Altitude T.P. (m)	pH	NaCl (%)	ORP (mV)	Organic matter ng/g-dry sediment)	Median particle size (μm)	Mud content (%)
May	Shirakawa	High	+1.494	N.M.	N.M.	210	6.3	143.4	31.8
		Low	-0.476	7.78	1.97	180	23.7	99.4	47.6
	Midorikawa	High	+0.553	7.68	0.76	-72	69.6	20.3	85.1
		Low	-0.833	8.60	1.34	-40	98.1	11.6	98.9
Jan	Shirakawa	High	N.M.	N.M.	N.M.	231	12.4	76.1	49.7
		Low	N.M.	7.55	0.60	227	14.7	40.9	59.4
	Midorikawa	High	N.M.	7.81	0.30	-113	47.2	42.9	67.8
		Low	N.M.	7.54	0.33	-21	76.5	17.9	86.0

T.P.: Tokyo peil, N.M.: Not measured

Homology analysis among bacterial DNA sequences

Partial 16S sequence data of the obtained clones analyzed by BLAST and their close phylum or class are summarized in Table 2. Many groups of bacteria were observed in the environmentally conserved Midorikawa and Shirakawa estuaries. In general, the dominant group was *Gamma-proteobacteria*, comprising 22.6% of the total clones, and the second most dominant group was *Alphaproteobacteria*, comprising 15.9%. *Chloroflexi*, *Deltaproteobacteria*, *Actinobacteria*, and *Bacteroidetes* were also common. The

microbial community composition among different sampling sites, seasons, and tidal positions are compared in Fig. 3. High ratios of *Deltaproteobacteria* and *Gammaproteobacteria* were detected in Midorikawa estuary compared to Shirakawa estuary, which seemed related to the slightly more anaerobic conditions with higher amounts of organic matter. However, *Gammaproteobacteria* in the sediment of Shirakawa-river estuary in May was not so high, the low value of organic matter was observed at both hightide and lowtide positions, as well.

The ratios of *Alphaproteobacteria*, *Actinobacteria*, and *Bacteroidetes* were higher in May than in January, possibly due to greater phototrophic production of organic compounds and subsequent aerobic degradation. In January, the ratio of *Gammaproteobacteria* at the low tide position was higher than that at the high tide position. While in May, the ratio of *Gamma*-*proteobacteria* at the high tide position was

higher than that at the low tide position. Although, the organic matter values at low tide positions were always higher than those at high tide positions.

Additionally, many *Gammaproteobacteria* clones were detected at the low tide position in the Midorikawa estuary, though they were uncultured bacteria.

Table 2. Closely related phylum or class for the detected clones as determined by homology analysis

Sampled season	May				January			
Place of estuary	Midorikawa		Shirakawa		Midorikawa		Shirakawa	
Tide level position	High	Low	High	Low	High	Low	High	Low
<i>Deltaproteobacteria</i>	3	3	1	0	7	4	2	2
<i>Firmicutes</i>	1	1	0	1	1	2	2	0
<i>Chloroflexi</i>	1	3	5	3	6	1	2	4
<i>Epsilonproteobacteria</i>	0	0	0	0	0	1	1	0
<i>Alphaproteobacteria</i>	4	2	4	10	7	4	4	3
<i>Betaproteobacteria</i>	0	0	0	0	0	1	1	1
<i>Gammaproteobacteria</i>	10	6	4	1	4	18	2	9
<i>Bacteroidetes</i>	2	6	5	2	1	1	2	1
<i>Actinobacteria</i>	1	0	4	7	3	1	6	1
<i>Acidobacteria</i>	1	2	1	0	6	2	4	4
<i>Nitrospirae</i>	0	2	0	1	0	1	1	1
<i>Planctomycetes</i>	1	0	2	1	2	0	0	2
<i>Verrucomicrobia</i>	1	0	1	0	0	0	0	1
<i>Cyanobacteria</i>	2	0	0	0	1	1	1	0
<i>Thermotogae</i>	0	0	0	2	0	0	0	0
TOTAL	27	25	27	28	38	37	28	29

1

Figure 3. Comparison of microbial communities in different sediment compositions (muddy, Midorikawa; sandy, Shirakawa), at different seasons, and at different tidal positions. Patterns indicating clones: solid, *Deltaproteobacteria*; dotted, *Firmicutes*; right-down diagonals, *Chloroflexi*; vertical stripes, *Epsilonproteobacteria*; wavy lines, *Alphaproteobacteria*; hatching, *Betaproteobacteria*; open, *Gammaproteobacteria*; stitched, *Bacteroidetes*; left-down diagonals, *Actinobacteria*; horizontal stripes, *Acidobacteria*; checkmarks, *Nitrospirae*; rectangular blocks, *Verrucomicrobia*; diamonds, *Cyanobacteria*; dashes, *Thermotogae*.

Figure 5. Neighbor-joining tree of *Alphaproteobacteria* clones based on partial 16S rRNA gene sequences

Figure 6. Neighbor-joining tree of *Deltaproteobacteria* clones based on partial 16S rRNA gene sequences

The SRB communities in the Midorikawa and Shirakawa estuaries were quite diverse. However, SRB were detected mainly in the Midorikawa estuary and were observed to a greater extent in January and at the high tide position. About 10% of the analyzed clones were SRB (Fig. 3), but few SOB clones were detected (Fig. 4), even though in an environmentally conserved tidal flat the hydrogen sulfide generated by SRB would likely be oxidized by SOB.

Anaerobic, facultatively anaerobic, and aerobic bacteria were observed in not just *Proteobacteria*, but also in the *Chloroflexi* group. However, in *Actinobacteria* and *Bacteroidetes* only aerobic bacteria were observed in both estuaries, both seasons, and both tidal positions (data not shown). Many of the identified clones were similar to salt-tolerant genera such as *Ilumatobacter* [22] and *Aestuariicola* [23]. The results presented here suggest that not only aerobic SOB, but also anaerobic phototrophic purple bacteria and anaerobic SRB might coexist in the top 5 cm of the sediments in the Shirakawa and Midorikawa.

At those conserved tidal flats, the ratio of *Gammaproteobacteria* and *Alphaproteobacteria* were higher than that of *Deltaproteobacteria*, the hydrogen sulfide producer. In comparison, at another location in Japan which is functionalized for intensive shellfish aquaculture, *Deltaproteobacteria* were detected as the most abundant [24]. Abundance of *Deltaproteobacteria* is stimulated by high content of organic matter and thus leads to enhancement of hydrogen sulfide production by the bacteria.

CONCLUSION

According to 16S rRNA, the bacterial community was found in high diversity with many clones close to uncultured strains in the tidal flats. Complex material cycles involving aerobic and anaerobic activities may mostly exist in the surface layer of sediment which was mainly performed by the abundant *Gammaproteobacteria*. The dominance of *Gammaproteobacteria* over anaerobic *Deltaproteobacteria* can be an indicator for unspoiled tidal flats.

REFERENCES

- [1] Sorokin, D.Y., T.P. Tourova, E.Y. Bezsoudnova, A. Pol, G. Muyzer. 2007. Denitrification in a binary culture and thiocyanate metabolism in *Thiohalophilus thiocyanoxidans* gen. nov. sp. nov. – a moderately halophilic chemolithoautotrophic sulfur-oxidizing Gammaproteobacterium from hypersaline lakes. *Arch. Microbiol.* 187. 441–450.
- [2] Andrade, L.L., D.C. Leite, E.M. Ferreira, L.Q. Ferreira, G.R. Paula, M.J. Maguire, C.R. Hubert, R.S. Peixoto, R.M. Domingues, A.S. Rosado. 2012. Microbial diversity and anaerobic hydrocarbon degradation potential in an oil-contaminated mangrove sediment. *Microbiol.* 12. 186-195.
- [3] Alejandro, A.G., R.M. Ramon, M. Silvia. 2013. Characterization of the anaerobic microbial community in oil-polluted subtidal sediments: aromatic biodegradation potential after the *Prestige* oil spill. *Environ. Microbiol.* 15. 77–92.
- [4] Jin, H.M., J.M. Kim, H.J. Lee, E.L. Madsen, C.O. Jeon. 2012. *Alteromonas* as a key agent of polycyclic aromatic hydrocarbon biodegradation in crude oil-contaminated coastal sediment. *Environ. Sci. Technol.* 46. 7731–7740.
- [5] Gillan, D.C., B. Danis, P. Pernet, G. Joly, P. Dubois. 2005. Structure of sediment-associated microbial communities along a heavy-metal contamination gradient in the marine environment. *Appl. Environ. Microbiol.* 71. 679–690.
- [6] Sun, M.Y., K.A. Dafforn, M.V. Brown, E.L. Johnston. 2012. Bacterial communities are sensitive indicators of contaminant stress. *Marine Poll. Bull.* 64. 1029-1038.
- [7] Tanaka, S., Y. Tashiro, N. Mitsutake, Y. Nakazono, G. Kobayashi, T. Kato, K. Kanda. 2011. Analysis of bacterial community structures in coastal sediments in Ariake sea. *Seibutsukogaku.* 89. 161-169.
- [8] Kim, B.K., Y.D. Park, H.M. Oh, J. Chun. 2009. Identification and characterization of metagenomic fragments from tidal flat sediment. *J. Microbiol.* 47. 402-410.
- [9] Mount, D.W. 2007. Using the Basic Local Alignment Search Tool (BLAST). Cold Spring Harb Protoc.
- [10] Thompson, J.D., T.J. Gibson, F. Plewniak, F. Jeanmougin, D.G. Higgins. 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 24. 4876-4882.
- [11] Tamura, K., J. Dudley, M. Nei, S. Kumar. 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.* 24. 1596-1599.

- [12] Japanese Industrial Standards Committee: Organic matter. In: Namiki, H. (Ed). Testing methods for industrial wastewater, JIS K 0102. Japanese Standards Association.
- [13] Yanagibayashi, M., Y. Nogi, L. Li, C. Kato. 1999. Changes in the microbial community in Japan Trench sediment from a depth of 6292 m during cultivation without decompression. *FEMS Microbiol. Lett.* 170. 271-279.
- [14] Yoon, J.H., T.K. Oh, Y.H. Park. 2004. *Kangiella koreensis* gen. nov., sp. nov. and *Kangiella aquimarina* sp. nov., isolated from a tidal flat of the Yellow Sea in Korea. *Int. J. Sys. Evol. Microbiol.* 54. 1829-1835.
- [15] Romanenko, L.A., N. Tanaka, G.M. Frolova, V.V. Mikhailov. 2010. *Kangiella japonica* sp. nov., isolated from a marine environment. *Int. J. Sys. Evol. Microbiol.* 60. 2583-2586.
- [16] Hozuki, T., T. Ohtsuka, K. Arai, K. Yoshimatsu, S. Tanaka, T. Fujiwara, 2010. Effect of salinity on hydroxylamine oxidation in a marine ammonia-oxidizing gamma-proteobacterium, *Nitrosococcus oceani* strain NS58: Molecular and catalytic properties of tetraheme cytochrome c-554. *Microbes Environ.* 25. 95-102.
- [17] Laurent, U., I. Laurent, L. Franc, L. Philippe. 2009. *Haliea rubra* sp. nov., a member of the *Gammaproteobacteria* from the Mediterranean Sea. *Int. J. Sys. Evol. Microbiol.* 59. 1188-1192.
- [18] Suzuki, T., Nakamura, T., and Fuse, H. 2012. Isolation of two novel marine ethylene-assimilating bacteria, *Haliea* species ETY-M and ETY-NAG, containing particulate methane monooxygenase-like genes, *Microbes Environ.* 27. 54-60.
- [19] Sorokin, D.Y., T.P. Tourova, G. Muyzer, G.J. Kuenen. 2008. *Thiohalospira halophila* gen. nov., sp. nov. and *Thiohalospira alkaliphila* sp. nov., novel obligately chemolitho-autotrophic, halophilic, sulfur-oxidizing gammaproteobacteria from hypersaline habitats. *Int. J. Sys. Evol. Microbiol.* 58. 1685-1692.
- [20] Daly, K., R.J. Sharp, A.J. McCarthy. 2000. Development of oligonucleotide probes and PCR primers for detecting phylogenetic subgroups of sulfate-reducing bacteria. *Microbiol.* 146. 1693-1705.
- [21] Suzuki, D., A. Ueki, A. Amaishi, K. Ueki. 2007. Diversity of substrate utilization and growth characteristics of sulfate-reducing bacteria isolated from estuarine sediment in Japan. *J. Gen. Appl. Microbiol.* 53. 119-132.
- [22] Matsumoto, A., H. Kasai, Y. Matsuo, S. Omura, Y. Shizuri, Y. Takahashi. 2009. *Ilumatobacter fluminis* gen. nov., sp. nov., a novel actinobacterium isolated from the sediment of an estuary. *J. Gen. Appl. Microbiol.* 55. 201-205.
- [23] Yoon, J.H., S.J. Kang, Y.T. Jung, T.K. Oh. 2008. *Aestuariicola saemankumensis* gen. nov., sp. nov., a member of the family *Flavobacteriaceae*, isolated from tidal flat sediment. *Int. J. Sys. Evol. Microbiol.* 58. 2126-2131.
- [24] Asami, H., M. Aida, K. Watanabe. 2005. Accelerated Sulfur Cycle in Coastal Marine Sediment beneath Areas of Intensive Shellfish Aquaculture. *Appl. Env. Microbiol.* 71. 2925-2933.